

Transverse Energy Measurement in $\sqrt{s_{NN}} = 62.4$ GeV Au+Au Collisions at RHIC

Raghunath Sahoo*,¹ Subhasis Chattopadhyaya,³ Alexandre A. P. Suaide,² Marcia Maria de Moura,² and D.P. Mahapatra¹
(for the STAR Collaboration)

¹*Institute of Physics, Bhubaneswar, India*

²*University of Sao Paulo, Brazil*

³*Variable Energy Cyclotron Centre, Kolkata, India*

The transverse energy distributions (E_T) have been measured for Au + Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV by the STAR experiment at RHIC. They have been obtained from two measurements, the hadronic transverse energy (E_T^{had}) and the electromagnetic transverse energy (E_T^{em}). E_T^{had} has been measured from the tracks obtained by Time Projection Chamber (TPC) excluding the electrons and positrons. E_T^{em} has been obtained by the STAR Barrel Electromagnetic Calorimeter (BEMC) which measures the energy of electrons, positrons and photons. The measure of transverse energy gives an estimate of the energy density of the fireball produced in heavy ion collisions. E_T per participant pair gives information about the production mechanism of particles.

PACS numbers:

Keywords: Transverse energy, Quark Gluon Plasma, Energy density

1. INTRODUCTION

The quest for understanding of the possible formation and existence of the quark-gluon plasma (QGP), the deconfined phase of quarks and gluons, has been a major area of heavy-ion research during the last couple of decades. The study of high energy nuclear collisions at the Relativistic Heavy Ion Collider (RHIC)[1] has opened a new domain in the exploration of strongly interacting matter at very high energy density. High temperature and densities may be generated in the most central nuclear collisions at RHIC, creating conditions in which a phase of deconfined quarks and gluons may exist [2,3]. The understanding of the early phase of the fireball produced in nuclear collisions requires the study of observables like the energy and momentum, produced transversely to the beam direction viz E_T and p_T . The number of charge particles produced and E_T are closely related to the collision geometry and are of importance in understanding the global properties of the system formed during the collision. Scattering of the partonic constituents of the incoming nuclei in the initial phase added to the rescattering of the produced partons and hadrons results in generation of E_T [4,5]. If the fireball of produced quanta breaks apart quickly without significant reinteraction, the observed dE_T/dy will be the same as that produced by the initial scattering. However, with interaction among the produced quanta, the system can achieve equilibrium at a very early state, after which it can expand resulting in a lowering of dE_T/dy [6,7]. To some extent this will be compensated by any transverse hydrodynamic flow[8]. Gluon saturation can delay the onset of the above hydrodynamic flow reducing the effective pressure and thereby reducing the difference in the initially produced and the observed E_T [9].

Here, we present preliminary results on E_T , produced in 62.4 GeV Au+Au collisions at RHIC. We present both the hadronic and the electromagnetic components of E_T , which were measured independently by the STAR detector using the Time projection Chamber (TPC) and the Barrel Electromagnetic Calorimeter (BEMC). A brief description of the STAR experiment with the detectors used together with the data analysis methods employed is given in Section-2. In Section-3 we present the results (STAR Preliminary) with conclusion presented in Section-4.

2. STAR EXPERIMENT AND DATA ANALYSIS

STAR[10], is an azimuthally symmetric, large acceptance solenoidal detector comprising of several detector subsystems. The subsystems relevant for this analysis are a large TPC located inside a 0.5 T solenoidal magnet, the BEMC

¹ * Corresponding author: Raghunath Sahoo¹ (email: raghu@iopb.res.in)

and two zero-degree calorimeters (ZDCs) for event selection. The BEMC[11] is a lead-scintillator sampling electromagnetic calorimeter with equal volumes of lead and scintillator. The full Barrel corresponds to 120 modules. Each module is composed of 40 towers (20 towers in η by 2 towers in ϕ), constructed to project to the center of the STAR detector. For the 2004 run, only half of the Barrel was instrumented, corresponding to a pseudorapidity coverage of $0 < \eta < 1$ with full azimuthal symmetry. The transverse dimensions of a tower are approximately $10 \times 10 \text{ cm}^2$, which at the radius of the front face of the detector corresponds to a phase space interval $(\Delta\eta, \Delta\phi)$ of $(0.05, 0.05)$. Each tower has a depth of 21 radiation lengths(X_0) corresponding to one interaction length for a hadron. The BEMC with a radius of 2.3 m sits inside the STAR solenoidal magnet. The electromagnetic energy resolution of the BEMC is $\delta E/E \sim 16\%/\sqrt{E(\text{GeV})}$.

The TPC[12] is the primary STAR detector used for the event reconstruction. It is a gas chamber, 4.2 m long with inner and outer radii of 50 and 200 cm respectively, placed in an uniform magnetic field of 0.5 T. The particles passing through the active gas medium release secondary electrons that drift to the readout end caps at both ends of the chamber. The readout system is based on multiwire proportional counters, with readout pads. There are 45 pad rows between the inner and outer radii of the TPC. The induced charge from the electrons is shared over several adjacent pads. The TPC provides up to 45 independent spatial and specific ionization dE/dx measurements. The dE/dx measurements, along with the momentum measurement from the bending of the tracks inside the magnetic field, determine the particle mass within limited kinematic regions. The TPC covers a pseudorapidity region with $|\eta| < 1.8$ with full azimuthal coverage.

The event trigger consisted of the coincidence of signals from two ZDCs, located at $\theta < 2 \text{ mrad}$ about the beam down stream of the first accelerator dipole magnet and are sensitive to spectator neutrons. These calorimeters provide a minimum bias trigger which, after collision vertex reconstruction, corresponds to $97 \pm 3\%$ of the geometric cross section σ_{geom}^{Au+Au} . The events are analyzed in centrality bins based on the TPC charged particle multiplicity in $|\eta| < 0.5$.

The present analysis is based on the minimum bias Au+Au collisions data at $\sqrt{s_{NN}} = 62.4 \text{ GeV}$, taken by STAR in the 2004 RHIC run. Here the TPC acceptance is limited by the BEMC. The TPC track quality cuts include a) z-coordinate (longitudinal axis) selection of collision vertex within 30 cm of the TPC center and b) a minimum TPC track space point cut of 10.

2.1. Hadronic Transverse Energy, E_T^{had}

The hadronic part of the transverse energy as measured from the momentum analysed TPC tracks is defined as

$$E_T^{had} = \sum_{tracks} E_{track} \sin\theta, \quad (1)$$

where the sum runs for all hadrons produced in the collision, except π^0 , η and other long-lived neutral hadrons. θ is the polar angle with respect to the beam axis and the collision vertex position. The Hadronic energy, E_{track} , is defined as [13,14]

$$E_{track} = \begin{cases} \sqrt{p^2 + m^2} - m, & \text{for nucleons} \\ \sqrt{p^2 + m^2} + m, & \text{for anti-nucleons} \\ \sqrt{p^2 + m^2}, & \text{for all other particles} \end{cases}$$

With the above definition,

$$E_T^{had} = C_0 \sum_{tracks} C_1(ID, p) E_{track}(ID, p) \sin\theta \quad (2)$$

where the sum includes all the primary tracks in BEMC acceptance. Here, C_0 is a correction factor defined as

$$C_0 = \frac{1}{f_{acc}} \frac{1}{f_{pT^{cut}}} \frac{1}{f_{neutral}} \quad (3)$$

where f_{acc} is the acceptance correction, $f_{neutral}$ is the correction for long-lived neutral hadrons not measured by the TPC, $f_{pT^{cut}}$ corresponding to the TPC low momentum cutoff. The factor $C_1(ID, p)$ is defined as

$$C_1(ID, p) = f_{bg}(pT) \frac{1}{f_{notID}} \frac{1}{eff(pT)} \quad (4)$$

which includes the uncertainty in the particle ID determination, f_{notID} , the momentum dependent tracking efficiency, $eff(pT)$ and the momentum dependent background, $f_{bg}(pT)$. Details of the procedure for finding out the correction factors are given elsewhere[13].

2.2. Electromagnetic Transverse Energy, E_T^{em}

E_T^{em} is measured from the BEMC tower hits corrected for the hadronic contaminations in the calorimeter. It is defined as

$$E_T^{em} = \sum_{towers} E_{tower}^{em} \sin(\theta_{tower}) \quad (5)$$

where E_{tower}^{em} is the electromagnetic energy measure in an BEMC tower and θ_{tower} is the polar angle of the center of the tower relative to the beam axis and the collision vertex position. Experimentally, E_T^{em} is given by

$$E_T^{em} = \frac{1}{f_{acc}} \sum_{towers} (E_{tower} - \Delta E_{tower}^{had}) \sin(\theta_{tower}) \quad (6)$$

where, f_{acc} is the acceptance correction, E_{tower} is the energy measured by an BEMC tower and ΔE_{tower}^{had} is the total correction for each tower to exclude the hadronic contributions. ΔE_{tower}^{had} is given by

$$\Delta E_{tower}^{had} = \frac{1}{f_{neutral}} \sum_{tracks} \frac{f_{elec}(pT)}{eff(pT)} \Delta E(p, \eta, d), \quad (7)$$

where, $\Delta E(p, \eta, d)$ is the energy deposited by a track projected on a BEMC tower as a function of it's momentum p , pseudorapidity η and distance d from the center of the tower to the track hit point. $f_{elec}(pT)$ is the correction to exclude electrons that are misidentified as hadrons. $eff(pT)$ is the track reconstruction efficiency and $f_{neutral}$ is the contribution to exclude the long-lived neutral hadron contribution.

3. RESULTS

Assuming the correction factors used in this analysis, to be the same as those obtained for 200 GeV Au+Au collisions[13], we have determined the minimum bias distributions of the total tranverse energy, E_T and it's components (E_T^{had} and E_T^{em}) separately. These are shown in Fig.1.(a)-(c)




FIG. 1: Minimum bias distributions of (a) total transverse energy, (b) hadronic transverse energy and (c) electromagnetic transverse energy, for $\sqrt{s_{NN}} = 62.4$ GeV Au+Au collisions.

In Fig. 2(a), we present dE_T/dy per participant pair for the top 5% central collisions, together with results from other experiments from AGS to RHIC[15-18]. The dE_T/dy values for this analysis, were calculated from $dE_T/d\eta$ using a factor of 1.18 obtained from HIJING simulation to convert η to y phase space. We obtained a value of 3.05 ± 0.05 (stat) GeV, consistent with an overall logarithmic growth of $dE_T/dy/(0.5N_{part})$ with $\sqrt{s_{NN}}$. For the same top 5% central events, $\langle E_T \rangle_{5\%}$ as obtained from $\langle dE_T/d\eta|_{\eta=0.5} \rangle$ for full azimuthal coverage and one unit η interval, is found to be 450 ± 6 (stat) GeV. Further, we have also determined the spatial energy density produced in the collision using the Bjorken formula[19]

$$\epsilon_{Bj} = \frac{dE_T}{dy} \frac{1}{\tau_0 \pi R^2} \quad (8)$$

where, dE_T/dy is the primordial rapidity density of transverse energy, R is the transverse system size given by $R = R_0 A^{\frac{1}{3}}$ and τ_0 is the formation time. Assuming $\tau_0 = 1$ fm/c, we get $\epsilon_{Bj} = 3.46 \pm 0.05$ GeV/fm³. This energy density is significantly higher than the energy density ~ 1 GeV/fm³ as required for the transition to a deconfined quark gluon plasma, predicted by lattice QCD [20]. This estimate is based on the assumption that local equilibrium has been achieved at $\tau \sim 1$ fm/c and then the system expands hydrodynamically.




FIG. 2: (a) $(dE_T/dy)/(0.5N_{part})$ Vs $\sqrt{s_{NN}}$ for central events, (b) Collision energy dependence of electromagnetic fraction of total E_T as given by $\langle dE_T^{em}/d\eta \rangle / \langle dE_T/d\eta \rangle$ for a number of systems from SPS to RHIC for central events

In Fig. 2(b), we have shown the electromagnetic fraction of the total transverse energy for the top 5% central events, as a function of the center of mass energy from SPS[21,22] to RHIC. This is seen to increase slowly when we go from AGS to RHIC. At 62.4 GeV this value is 0.29 ± 0.01 (stat). As discussed in ref.[13], the observed electromagnetic fraction of the total transverse energy is strongly dependent on the baryon to meson ratio. At very high energy it is expected that virtually all the E_T will be carried by mesons, as an almost baryon free region is expected to be created in the central rapidity region, while at lower SPS energies, baryon dominance results in a much smaller electromagnetic fraction. A very high value of the electromagnetic fraction of total transverse energy is expected in case of a long-lived deconfined phase, due to an excess yield of photons[23]. Based on the present ratio of 0.29 ± 0.01 (stat) it is very difficult to conclude anything regarding the formation of a deconfined QGP phase.

4. CONCLUSIONS

In the present work we have reported preliminary STAR results on E_T within $0 < \eta < 1$ for Au+Au collisions at $\sqrt{s_{NN}} = 62.4$ GeV. For top 5% central events $\langle E_T \rangle$ has been estimated to be 450 ± 6 (stat) GeV. The value of $\langle dE_T/dy \rangle / (0.5N_{part})$ has been found to be 3.05 ± 0.05 (stat) GeV. Knowing that the observed E_T is lower than the initial values [6,9,24], the present value of 3.05 ± 0.05 (stat) GeV may be considered as a lower bound only. The initial energy density estimated within the framework of boost-invariant hydrodynamics, as given by ϵ_{Bj} , has been found to be 3.46 ± 0.05 (stat) GeV/fm³ which is well above that required for the deconfinement phase transition as predicted by lattice QCD[20]. The electromagnetic fraction of the total transverse energy for central events is found to be 0.29 ± 0.01 (stat), consistent with the fact that the final state is dominated by mesons. Finally, this method of independent measurement of E_T^{em} and E_T^{had} gives an unique opportunity to study event by event fluctuations in these observables and in their ratios. The correction factors and the systematic errors are yet to be calculated for this analysis.

Acknowledgments

We thank the RHIC Operations Group and RCF at BNL, and the NERSC Center at LBNL for their support. This work was supported in part by the HENP Divisions of the Office of Science of the U.S. DOE; the U.S. NSF; the BMBF of Germany; IN2P3, RA, RPL, and EMN of France; EPSRC of the United Kingdom; FAPESP of Brazil; the Russian Ministry of Science and Technology; the Ministry of Education and the NNSFC of China; IRP and GA of the Czech Republic, FOM of the Netherlands, DAE, DST, and CSIR of the Government of India; Swiss NSF; the

Polish State Committee for Scientific Research; STAA of Slovakia, and the Korea Sci. & Eng. Foundation.

-
- [1] T. Roser, *Nucl. Phys.* **A698**, 23c(2002).
 - [2] J. P. Blaizot, *Nucl. Phys.* **A661**, 3c (1999).
 - [3] P. Jacobs and X.N. Wang, hep-ph/0450125.
 - [4] M. Jacob and P.V. Landshoff, *Mod. Phys. Lett.* **A1**, 657 (1986)
 - [5] X. N. Wang, *Phys. Rep.* **280**, 287 (1997)
 - [6] M. Gyulassy and T. Matsui, *Phys. Rev. D* **29**, 419 (1984)
 - [7] K. J. Eskola et al., *Nucl. Phys.* **B570**, 379 (2000)
 - [8] P. F. kolb et al., *Nucl. Phys.* **A696**, 197 (2001)
 - [9] A. Dumitru and M. Gyulassy, *Phys. Lett.* **B494**, 215 (2000)
 - [10] STAR Collaboration, K. H. Akermann et al., *Nucl. Instr. and Meth. Phys.* **A 499**, 624 (2003)
 - [11] M. Beddo et al., *Nucl. Instr. and Meth. Phys.* **A 499**, 725 (2003)
 - [12] M. Anderson et al., *Nucl. Instr. and Meth. Phys.* **A 499**, 659 (2003)
 - [13] STAR Collaboration, J. Adams et al., *Phys. Rev.* **C 70**, 054907 (2004)
 - [14] HELIOS Collaboration, T. Akesson et al., *Z. Phys.* **C38**, 383 (1988)
 - [15] J. Barrette et al., *Phys. Rev. Lett.* **70**, 2996 (1993)
 - [16] T. Alber et al., *Phys. Rev. Lett.* **75**, 3814 (1995)
 - [17] WA98 Collaboration, M. M. Aggarwal et al., *Eur. Phys. J.* **C18**, 651 (2001)
 - [18] PHENIX Collaboration, K. Adcox et al., *Phys. Rev. Lett.* **87**, 052301 (2001)
 - [19] J. D. Bjorken, *Phys. Rev. D* **27**, 140 (1983)
 - [20] F. Karsch, *Nucl. Phys.* **A698**, 199c (2002)
 - [21] WA80 Collaboration, R. Albrecht et al., *Phys. Rev.* **C 44**, 2736 (1991)
 - [22] WA98 Collaboration, Systematics of Transverse Energy Production in Pb induced Collisions at 158 GeV/nucleon (1996)
 - [23] F. Halzen and H. C. Liu, *Phys. Rev. D* **25**, 1842 (1982)
 - [24] PHENIX Collaboration, K. Adcox et al., *Nucl. Phys.* **A757**, 102 (2005)